

Name.	Distribution of the Order, Genus, or Species.
COMPOSITEÆ.	
* <i>Lagenophora</i>	Australasia to India and the Sandwich Islands, with a few species in South America, one of which reaches Tristan da Cunha.
* <i>Abrotanella</i>	Several species in Australasia and one in the Falklands.
<i>Cotula plumosa</i>	Crozets, Kerguelen, Macquarie, Auckland, and Campbell Islands.
<i>Flaveria</i>	Seven species, six of which inhabit America, ranging from Chili to Mexico and Florida, and one Australia, the Australian species being very near one of the commoner American ones.
* <i>Microseris</i>	A genus of about a dozen species, whereof one is found in Extratropical South America, one in Australasia, and the rest in North-west America.
STYLIDÆÆ.	
* <i>Phyllachne</i>	Australasia and Fuegia.
LOBELIACEÆ.	
* <i>Lobelia anceps</i>	Australasia, South America, and South Africa.
* <i>Pratia</i>	Endemic species in Australasia, South America, and Asia.
* <i>Selliera radicans</i>	Australasia and Chili.
ERICACEÆ.	
* <i>Pernettya</i>	One species endemic in Tasmania and New Zealand, and the rest in South America.
*ERICACEÆ, sub-ord. EPACRIDÆÆ	Restricted to Australasia, Polynesia, Malayan Peninsula and Archipelago, except the monotypic <i>Lebetanthus americanus</i> , and specially numerous in Australia.
PRIMULACEÆ.	
* <i>Samolus repens</i>	Australasia, Polynesia, and South Chili.
GENTIANACEÆ.	
* <i>Gentiana saxosa</i>	The only species found in Australia, with which Mueller unites the South American <i>Gentiana magellanica</i> .
CONVOLVULACEÆ.	
<i>Calystegia tuguriorum</i>	New Zealand, Masafuera, Chili.
SCROPHULARINEÆ.	
<i>Calceolaria</i>	Two endemic species in New Zealand, and many in America, ranging from the Falklands and Fuegia to Mexico.
* <i>Gratiola peruviana</i>	Australasia and South America.
<i>Veronica elliptica</i>	Australasia, Falklands, and Fuegia.
<i>Ourisia</i>	One species endemic in Tasmania, about six in New Zealand, and about a dozen in America, ranging from Fuegia to New Grenada.
PLANTAGINEÆ.	
* <i>Plantago</i>	<i>Plantago brownii</i> , found in Tasmania and New Zealand, has a representative species in the South American <i>Plantago barbata</i> .
* <i>Euphrasia antarctica</i>	Australasia and Chili.
POLYGONEÆ.	
* <i>Muehlenbeckia</i>	Endemic species in Australasia and in the Andes.