

Dicranella condensata, J. Angstr.

Dicranella condensata, J. Angstr., in *Hedwigia*, 1875, p. 92.

ST HELENA.

The description of this seems to indicate *Dicranella ascensionica*, Mitt.; but this species is absent in all our collections from St Helena.

Campylopus introflexus, Hedw.

Campylopus introflexus, Hedw., Sp. Musc., t. 20 (*Dicranum*); Mitt. in Melliss's St Hel., p. 358.

ST HELENA.—Damp banks on the high land—*Burchell*; *Melliss*.

A common moss both in the northern and southern hemispheres.

Campylopus helenicus, C. Müll.

Campylopus helenicus, C. Müll, Synop. Musc. Frond., ii. p. 599 (*Dicranum*); Mitt. in Melliss's St Hel., p. 358.

ST HELENA.—Endemic.

One of the commoner species, figuring in every collection.

Campylopus scabricuspis, Mitt.

Campylopus scabricuspis, Mitt. in Melliss's St Hel., p. 358.

ST HELENA.—Endemic. Damp roadside banks on the high land—*Melliss*.

Glyphomitrium helenicum, Mitt.

Glyphomitrium helenicum, Mitt. in Melliss's St Hel., p. 359.

ST HELENA.—Endemic.

Only collected by Captain Haughton.

Ceratodon purpureus, Linn.

Ceratodon purpureus, Linn. (*Mnium*); C. Müll, Synop. Musc. Frond., i. p. 646; Melliss, St Hel., p. 359.

ST HELENA. Damp banks on the high land—*Burchell*; *Melliss*.

The most widely diffused of all mosses.

Tortula crispula, Bruch.

Tortula crispula, Bruch. (*Trichostomum*); C. Müll, Synop. Musc. Frond., i. p. 571; Melliss, St Hel., p. 360.

ST HELENA.—Indigenous? Old walls and stones in the high land—*Melliss*.

Europe; North Africa.