

Thelotrema sp. ? Crombie.*Thelotrema* sp. ? Crombie in Journ. Linn. Soc. Lond., xvi. p. 223.JUAN FERNANDEZ. *Moseley*.

"On the bark of trees : indeterminable."

Urceolaria *scruposa*, Acharius.*Urceolaria scruposa*, Acharius, Lich. Univ., p. 338; Nyl., Lich. Scand., p. 176; Crombie in Journ. Linn. Soc. Lond., xvi. p. 223.JUAN FERNANDEZ. *Moseley*.

A common European species.

Lecidea *fernandezii*, Crombie.*Lecidea fernandezii*, Crombie in Journ. Linn. Soc. Lond., xvi. p. 223."Omnino est similis *Lecideæ urceolaria*, Nyl. in Flora, 1873, p. 298, differt autem hypothecio et perithecio fuscis, sporisque fere minoribus, longit. 0·010–0·014 millim., crassit. 0·004–0·006 millim. Forsan varietas *Lecideæ urceolaria*."JUAN FERNANDEZ.—Parasitic on *Urceolaria scruposa*—*Moseley*.

ALGÆ.

MELANOSPERMEÆ.

Zonaria *variegata*, Lamour.*Zonaria variegata*, Lamour., Dict., p. 11; J. G. Agardh, Sp. Alg., i. p. 108; Dickie in Journ. Linn. Soc. Lond., xv. p. 452.‡JUAN FERNANDEZ.—On *Corallina subulata*. *Moseley*.

Chiefly in the Atlantic; also in the Indian Ocean and on the coast of Australia.

Dictyota *kunthii*, J. G. Agardh.*Dictyota kunthii*, J. G. Agardh, Sp. Alg., i. p. 94, et Ic. ined., t. 15; Gay, Fl. Chil., viii. p. 260; Dickie in Journ. Linn. Soc. Lond., xv. p. 452; Hook. f., Handb. Fl. N. Zeal., p. 657.JUAN FERNANDEZ. *Moseley*.

Common on the coasts of Chili and New Zealand, and extending northward in the Pacific to California.

Dictyota *nævosa*, Suhr.*Dictyota nævosa*, Suhr; Eckel, n. 8; J. G. Agardh, Sp. Alg., i. p. 95; Dickie in Journ. Linn. Soc. Lond., xv. p. 452.JUAN FERNANDEZ. *Moseley*.

South Africa; Tasmania.