

**Citrullus vulgaris, Schrad.**

*Citrullus vulgaris*, Schrad. ; Hook. f., Fl. Brit. Ind., ii. p. 621 ; Seem., Fl. Vit., p. 104 ; Oliver, Fl. Trop. Afr., ii. p. 549.

*Citrullus edulis*, Spach. ; Miq., Fl. Ind. Bat., i. 1, p. 662.

*Cucurbita citrullus*, Linn. ; Jouan in Mém. Soc. Sci. Nat. Cherbourg, xi. 1865, p. 149.

MOA.—The Water-melon exists in a wild and cultivated state in nearly all tropical and many subtropical countries, but it is believed to be really indigenous only in Tropical Africa. It is cultivated, and more or less wild, in many of the islands of Polynesia, where, however, according to Seemann, it is known to have been introduced by Europeans.

**Momordica charantia, Linn.**

*Momordica charantia*, Linn. ; Miq., Fl. Ind. Bat., i. 1, p. 663 ; Hook. f., Fl. Brit. Ind., ii. p. 616 ; Oliver, Fl. Trop. Afr., ii. p. 537 ; Seem., Fl. Vit., p. 105.

TIMOR LAUT.—Commonly cultivated and naturalised in tropical countries, and probably indigenous in Tropical Africa. Seemann collected it in the Fijis, and he mentions that it was collected in Tahiti by Banks and Solander. *Momordica balsamina*, Linn., has nearly the same range as the present, and also extends to Australia. Both species are regarded by Cogniaux (DC., Monogr. Phanerog., iii. pp. 437 et 440) as introduced in America.

**Lagenaria vulgaris, Seringe.**

*Lagenaria vulgaris*, Seringe ; Benth., Fl. Austr., iii. p. 316 ; Miq., Fl. Ind. Bat., i. 1, p. 669 ; Jouan in Mém. Soc. Sci. Nat. Cherbourg, xi. 1865, p. 149 ; Seem., Fl. Vit., p. 106 ; Nadeaud, Enum. Pl. Tahiti, p. 66.

The Calabash Gourd, generally cultivated for the fruit, which is not eaten, but used for making bottles, &c. It was found in Easter Island, the Sandwich Islands, and other parts of Polynesia, when first discovered by Europeans.

**Cucumis trigonus, Roxb. ?**

*Cucumis trigonus*, Roxb. ? Hook. f., Fl. Brit. Ind., ii. p. 619 ; Benth., Fl. Austr., iii. p. 317 ; Miq., Fl. Ind. Bat., i. 1, p. 671 (*Cucumis turbinatus*).

TIMOR LAUT.—Generally diffused in Tropical Asia and extending to North Australia.

**Zehneria sp. (aff. *Zehneria mucronata*, Miq.)**

TIMOR LAUT.—The species may be different from the Java plant. *Zehneria*, as circumscribed by Bentham and Hooker, is confined to the Old World, with the exception of one species in Peru.