

Cerbera odollam, Gærtn.

Cerbera odollam, Gærtn. ; Benth., Fl. Austr., iv. p. 306 ; Miq., Fl. Ind. Bat., ii. p. 413 ; Hook. f., Fl. Brit. Ind., iii. p. 638.

Cerbera lactaria, Hamilt. ; Seem., Fl. Vit., p. 638.

ARROU ; KI ; MOA ; DIA.—A small tree inhabiting the coast region and salt-marshes of Tropical Asia, Australia, and many of the islands of Polynesia, including the New Hebrides, the Tongan and Fiji Islands, and the Isle of Pines, near New Caledonia. The large fibrous seed-vessel of this and the three or four other species of the genus, which are also sea-shore trees, floats on the water, and is diffused by oceanic currents. It is in the collection of drift fruits and seeds made by Mr Moseley.

Ochrosia borbonica, Gmel. ?

Ochrosia borbonica, Gmel. ? Hook. f., Fl. Brit. Ind., iii. p. 638.

ARROU.—Imperfect specimens of an *Ochrosia*, probably of this species, which inhabits the tidal forests of the Andaman Islands, Singapore, Java, Ceylon, the Seychelles and Mascarene Islands. Another species was collected in Timor Laut, perhaps the Australian *Ochrosia elliptica*, Labill. There are about a dozen species in the region indicated, some of them extending to the Pacific Islands. Darwin collected one species, *Ochrosia parviflora*, Henslow, in the Keeling Islands, and the same one has since been found in Tahiti, the Ellice, Samoan, and Fiji groups.

Plumeria acutifolia, Poir.

Plumeria acutifolia, Poir. ; Miq., Fl. Ind. Bat., ii. p. 426 ; Hook. f., Fl. Brit. Ind., iii. p. 641.

KI.—A tree of American origin, now commonly cultivated and widely naturalised in India and the Archipelago. The genus, which numbers from thirty to forty species, is wholly American, though several species are now colonised in the Old World.

Tabernæmontana parviflora, Dcne.

Tabernæmontana parviflora, Dcne., Herb. Timor. Descr., p. 51.

Tabernæmontana decaisnei, DC., Prodr., viii. p. 369 ; Miq., Fl. Ind. Bat., ii. p. 420.

KI ; LAKOR ; TIMOR LAUT ; DAMMAR ; MOA ; LETTI ; BABAR.—Bentham (Fl. Austr., iv. p. 311) regards this as a form of *Tabernæmontana orientalis*. There are upwards of 100 species of this genus, spread over the warmer regions of both hemispheres.

Tabernæmontana orientalis, R. Br. var.

Tabernæmontana orientalis, R. Br. var. ; Benth., Fl. Austr., iv. p. 311 ; Seem., Fl. Vit., p. 159, excl. syn. *Tabernæmontana cumingiana*, DC., *fide* Benth.

TIMOR LAUT.—A sea-coast tree or shrub found on the shores of New South Wales, Queensland, North Australia, the Fijis, and other Pacific Islands.