

**Arthonia polymorphoides**, Nyl.

*Arthonia polymorphoides*, Nyl. in Journ. Linn. Soc. Lond., xvi. p. 216; Crombie in Journ. Linn. Soc. Lond., xvi. p. 226.

*Arthonia interveniens*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 372, non Nyl.

ADMIRALTY ISLANDS.—On thin bark of trees: sparingly gathered. (Spores 0·016–0·022 millim. long, 0·006–0·007 millim. thick.)

This species was founded on Bermudan specimens collected by Mr Moseley, and these two distant localities are the only ones hitherto recorded for it.

**Platygrapha stenogramma**, Nyl.

*Platygrapha stenogramma*, Nyl. ex Crombie in Journ. Linn. Soc. Lond., xvi. p. 226.

Thallus albidus subopacus tenuis lævis, tenuiter nigro limitatus; apothecia nigra lineari-elongata (latit. circiter 0·15 millim., longit. sæpius 1–2 millim.), simplicia, plus minusve flexuosa, intus albida; sporæ 8-næ aciculari-fusiformes 7–9 septatæ, longit. 0·030–0·035 millim., crassit. 0·004 millim., paraphyses irregulares graciles, hypothecium incolor. Iodogelatina hymenialis dilute vinose rubescens.

Species apotheciis elongatis thallino-obductis (epithecio solo nudo) bene distincta, facie fere *Graphidis pavonianæ*. Spermatia arcuata, longit. 0·018–0·023 millim., crassit. 0·0005 millim.

ADMIRALTY ISLANDS.—On the bark of trees; probably not uncommon.

**Graphis scripta**, Acharius.

*Graphis scripta*, Acharius; Crombie in Journ. Linn. Soc. Lond., xvi. p. 226 (*forma pulverulenta*).

ADMIRALTY ISLANDS.—On the bark of trees; sufficiently typical.

This common British lichen is spread nearly all over the world.

**Graphis dendrogramma**, Nyl.

*Graphis dendrogramma*, Nyl. ex Crombie in Journ. Linn. Soc. Lond., xvi. p. 226.

Affinis *Graphidi scriptæ*, sed apothecia cæsio-pruinosa longiora dendroideo-divisa, thalodeo-marginata; sporæ 8–10 loculares, longit. 0·020–0·030 millim., crassit. 0·006–0·007 millim.

ADMIRALTY ISLANDS.—On the bark of trees; very sparingly gathered.

**Verrucaria paramera**, Nyl.

*Verrucaria paramera*, Nyl. ex Crombie in Journ. Linn. Soc. Lond., xvi. p. 227.

Thallus macula luteo-pallescente indicatus; apothecia pyrenio integre nigro, parte dimidia superiore nuda convexa prominula (latit. 0·5–0·7 millim.); sporæ 8-næ, fuscæ vel fuscescentes, ellipsoideæ vel oblongo-ellipsoideæ, 6-loculares vel demum oculis quibusdam