

Family, Genus, and Species.	Range in Depth.
<i>Cynthia cerebriformis</i> , . . . . .	6-15 fathoms.
<i>fissa</i> , . . . . .	38-40    "
<i>formosa</i> , . . . . .	3-11     "
<i>arenosa</i> , . . . . .	1-8      "
<i>irregularis</i> , . . . . .	2-10     "
<i>pallida</i> , . . . . .	10-20    "
<i>papictensis</i> , . . . . .	10-20    "
<i>complanata</i> , . . . . .	6         "
<i>hispida</i> , . . . . .	38-40    "
<i>Styela bythia</i> , . . . . .	2600     "
<i>squamosa</i> , . . . . .	2600     "
<i>grandis</i> , . . . . .	150      "
<i>gyrosa</i> , . . . . .	6         "
<i>convexa</i> , . . . . .	150      "
<i>lactea</i> , . . . . .	10-100   "
<i>exigua</i> , . . . . .	2-10     "
<i>clava</i> , . . . . .	8-50     "
<i>oblonga</i> , . . . . .	600      "
<i>flava</i> , . . . . .	600      "
<i>glans</i> , . . . . .	600      "
<i>radicosa</i> , . . . . .	38-40    "
<i>Bathyoncus mirabilis</i> , . . . . .	1600     "
<i>Polycarpa viridis</i> , . . . . .	2-15     "
<i>tinctor</i> , . . . . .	2-15     "
<i>minuta</i> , . . . . .	150      "
<i>molguloides</i> , . . . . .	38-40    "
<i>quadrata</i> , . . . . .	129      "
<i>pilella</i> , . . . . .	7-20     "
<i>rigida</i> , . . . . .	38-40    "
<i>longisiphonica</i> , . . . . .	6-15     "
<i>irregularis</i> , . . . . .	18        "
<i>sulcata</i> , . . . . .	17        "
<i>pedata</i> , . . . . .	10-20    "
<i>radicata</i> , . . . . .	6 and 120   "
<b>ASCIDIIDE—</b>	
<i>Corynascidia suhmi</i> , . . . . .	1375-2160   "
<i>Corella japonica</i> , . . . . .	8-50        "
<i>Abyssascidia wyvillii</i> , . . . . .	2600        "
<i>Pachychlæna obesa</i> , . . . . .	38-40       "
<i>gigantea</i> , . . . . .	10-20       "
<i>oblonga</i> , . . . . .	38-40       "
<i>Ascidia challengerii</i> , . . . . .	10-60       "
<i>vasculosa</i> , . . . . .	28          "
<i>placenta</i> , . . . . .	150         "
<i>nigra</i> , . . . . .	Shallow water, and 150 fathoms.