

## PLATE XVI.

Figs. 1–5. *Cynthia formosa*, Herdman.

Figs. 6–9. *Cynthia arenosa*, Herdman.

Figs. 10–12. *Cynthia irregularis*, Herdman.

Fig. 1. *Cynthia formosa*, from the right side ; natural size.

Fig. 2. Part of the branchial sac of *Cynthia formosa*, seen from the inside ; magnified (Hartn., obj. 4).

Fig. 3. A small portion of the branchial sac of *Cynthia formosa*, seen from the inside, showing an irregularity ; magnified (Hartn., obj. 4).

Fig. 4. Part of the dorsal lamina of *Cynthia formosa*, showing the marginal languets (*l.*) ; magnified (Hartn., obj. 4).

Fig. 5. The dorsal region of the anterior end of the branchial sac, showing the dorsal tubercle (*d.t.*), the peritubercular area, the præbranchial zone, and part of the circlet of tentacles (*tn.* and *tn'*.); magnified (Hartn., obj. 4).

Fig. 6. *Cynthia arenosa*, a specimen seen from the right side ; natural size.

Fig. 7. Part of the branchial sac of *Cynthia arenosa*, seen from the inside, and showing two of the folds (*br.f.*), two interspaces, and part of the dorsal lamina (*l.*) ; magnified (Hartn., obj. 4).

Fig. 8. The dorsal tubercle and peritubercular area of *Cynthia arenosa* ; magnified (Hartn., obj. 4).

Fig. 9. A small portion of the modified test lining the branchial siphon of *Cynthia arenosa* ; magnified (Hartn., obj. 5).

Fig. 10. *Cynthia irregularis*; from the left side ; natural size.

Fig. 11. Part of the branchial sac of *Cynthia irregularis*, seen from the inside, and showing a fold (*br.f.*), and several irregular areas ; magnified (Hartn., obj. 4).

Fig. 12. The peritubercular area and dorsal tubercle of *Cynthia irregularis* ; magnified (Hartn., obj. 4).