

PLATE X.

		Diam.	Page
Fig. 1. <i>Picroceroides tubularis</i> , n. gen. and sp., adult male,	. . . . . ×	1½	77
Fig. 1a. Lateral view (in outline) of the carapace,	. . . . . ×	1½	
Fig. 1b. Inferior view of the cervical region, showing the eyes, antennæ, and exterior maxillipedes,	. . . . . ×	2½	
Fig. 1c. Chelipede,	. . . . . ×	2½	
Fig. 2. <i>Macrocaeloma concava</i> , n. sp., adult male,	. . . . . ×	1½	81
Fig. 2a. Lateral view of carapace in outline,	. . . . .	nat. size.	
Fig. 2b. Inferior view of the cervical region, showing the form of the antennæ and exterior maxillipedes,	. . . . . ×	2	
Fig. 3. <i>Mithrax</i> sp., young male example,	. . . . . ×	5	89
Fig. 3a. Inferior view of the cervical region, showing the basal antennal joint and maxillipedes; considerably magnified.			
Fig. 3b. Chelipede of the same specimen.			
Fig. 4. <i>Lambrus intermedius</i> , Miers, adult male (from the type- specimen, from the Korean Channel, in the collection of the British Museum),	. . . . . × about 2		96
Fig. 5. <i>Lambrus (Aulacolambrus) hoplonotus</i> , var. <i>granulosus</i> , Miers, adult female,	. . . . . ×	2	98