

North Pacific.

<i>Sergestes præcollis.</i>		<i>Sergestes ventridentatus.</i>
<i>Sergestes spiniventralis.</i>		<i>Sergestes utrinquedens.</i>
<i>Sciacaris telsonis.</i>		

STATION 267. August 28, 1875; lat. 9° 28' N., long. 150° 49' W.; south of the Sandwich Islands. Tow-net at 2000 fathoms.

Gennadas parvus (1).

STATION 270. September 4, 1875; lat. 2° 34' N., long. 149° 9' W.; surface temperature, 79°·5. Surface.

<i>Lucifer typus.</i>		<i>Lucifer reynaudii.</i>
-----------------------	--	---------------------------

South Pacific, north of Low Archipelago; September 1875.

<i>Sergestes rinki.</i>		<i>Sergestes oculatus.</i>
<i>Caricyphus gibberosus.</i>		

STATION 272. September 8, 1875; lat. 3° 48' S., long. 152° 56' W.; north of the Low Archipelago; depth, 2600 fathoms; bottom, Radiolarian ooze; bottom temperature, 35°·1. Trawled.

Benthesicymus crenatus (1 ♀).

Between Honolulu and the Low Islands, September 12, 1875.

Sergestes armatus.

STATION 276. September 16, 1875; lat. 13° 28' S., long. 149° 30' W.; near the Low Archipelago; depth, 2350 fathoms; bottom, red clay; bottom temperature, 35°·1. Trawled.

<i>Aristeus armatus</i> (1 ♀).		<i>Benthesicymus crenatus</i> (2 ♀).
--------------------------------	--	--------------------------------------

STATION 279C. October 2, 1875; lat. 17° 29' 11" S., long. 149° 34' 32" W.; near Tahiti; depth, 680 fathoms; bottom, volcanic mud. Trawled.

Alpheus spiniger (1 ♀).

Tahiti.

<i>Panulirus penicillatus</i> (1).		<i>Bithynis lar</i> (5 ♂). Fresh-water.
------------------------------------	--	---