

PLATE XXVIII.

- | | PAGE |
|---|------|
| Figs. 1-11. <i>Hyalonema divergens</i> , F. E. S., | 199 |
| Fig. 1. Somewhat damaged specimen, preserved in alcohol ; natural size. | |
| Fig. 2. Marginal oxydiact with four central knobs ; $\times 100$. | |
| Fig. 3. Great amphidisc of the dermal membrane ; $\times 300$. | |
| Figs. 4, 5. Smaller amphidiscs ; $\times 300$. | |
| Fig. 6. Oxyhexact with smooth bent rays ; $\times 300$. | |
| Fig. 7. Autodermal pentact pinulus ; $\times 100$. | |
| Fig. 8. Oxyhexact with smooth straight rays ; $\times 300$. | |
| Fig. 9. Oxyhexact with smooth bent rays ; $\times 300$. | |
| Figs. 10, 11. Strong short spicules, triact and tetract, with rough ends, from the basal pad of the body ; $\times 100$. | |
| Figs. 12-17. <i>Hyalonema lusitanicum</i> , Bocage, | 225 |
| Fig. 12. Small amphidisc ; $\times 300$. | |
| Fig. 13. Middle-sized amphidisc ; $\times 300$. | |
| Fig. 14. Greater amphidisc with nodular axial beam ; $\times 300$. | |
| Fig. 15. Oxyhexact with smooth bent rays ; $\times 300$. | |
| Fig. 16. Oxyhexact with smooth straight rays ; $\times 300$. | |
| Fig. 17. Autodermal pentact pinulus ; $\times 300$. | |