

of ocelli conterminous, of nearly equal extent, the upper groups triangular, the lower almost circular.

*Upper Antennæ.*—Peduncle moderately stout; first joint of flagellum longer than the peduncle, set at right angles to it, carrying a large brush of filaments; second joint attached at the apex of the first, a little longer than broad, having some long apical filaments; the third joint considerably longer and more slender, with filaments some way from the apex; the fourth joint much shorter and thinner than the third, with some apical setules.

*Lower Antennæ.*—Third (first free) joint of the peduncle rather more than a quarter as long as the following joint, distally narrowed; fourth joint long and slender, distally widened; fifth joint equal in length to the fourth; first joint of flagellum slender, three-quarters of the length of the preceding joint; second joint rather more than half the length of the first.

*First Gnathopods.*—Side-plates with convex front margin. The first joint a little widened below; the second longer than broad; the third about three times as long as wide, longer and broader than the fourth joint, with a hair-like setule at each apex and one above the centre of the hind margin; the fourth joint similar, but a little narrowed distally, with a setule at each apex; the fifth joint shorter and narrower than the fourth, a little curved, distally narrowed; the finger little curved, acute, less than half the length of the fifth joint; the whole limb peræopod-like.

*Second Gnathopods* similar to the first, but scarcely so long, having the first and second joints a little longer, but the third and fourth decidedly shorter, each of the two latter joints having the apical setules and a setule near the middle of the hind margin; there is also a setule near the apex of the second joint.

*First Peræopods* like the gnathopods, but rather longer, the difference in length depending chiefly on the fourth and fifth joints which are about equal in length; there is a spinule near the hinder apex of the second joint and one high up on the hind margin of each of the three following joints, besides subapical setules.

*Second Peræopods* a little longer than the first.

*Third Peræopods.*—The side-plates nearly as deep as the breadth, the margins convex; within the side-plates there is a long straight spine-like process pointing backwards. The first joint broadly oval, almost smooth-edged, a little shorter than the third, fourth, and fifth joints together; the second joint scarcely longer than broad, with a setule above the front apex; the third joint very much shorter than the fourth, with three setules on the front margin; fourth joint longer than the fifth, with a setule at the apex behind, and three setules on the front margin, which is sparsely and finely pectinate; the fifth joint slender, tapering, longer than the third, having three setules on the hind margin and one on the front, which has a little scarcely perceptible pectination; the finger not a quarter the length of the preceding joint.