

Hyperia—continued.

	{ 83, 189, 194, 223, 274 282, 317, 321, 331, 337 358, 382, 421, 430, 465 494, 495, 548, 588, 592 593, 1625, 1640
Galba, Gould, 1841,	
{ Gaudichaudi, Bovallius, 1887,	588
Gaudichaudii, M.-Edw., 1840,	189, 232, 1394 , pl. clxix
gracilipes, Norman, 1869,	1420, 1628
Krocyeri, Bovallius, 1885,	46, 54, 179, 558
Latreillei, Gosse, 1855,	282, 588, 592, 593, 1645
latreilli, De Kay, 1844,	207
{ Latreillii, M.-Edw., 1830,	{ 139, 142, 163, 175, 189 194, 221, 223, 271, 306 321, 547, 1625
Lesueurii, M.-Edw., 1840,	106, 175, 189
luzoni, Stebbing, 1888,	1382 , pl. clxvi, A
macrocephala, Sp. Bate, 1862	264, 268
Martinezii, Fritz Müller, 1864,	349, 1399
mediterranea, Carus, 1885,	561, 588
medusarum, ¹ Bovallius, 1887,	42, 45 , 592, 593, 1617
Medusarum, Sp. Bate, 1862,	358, 382, 450, 1628
medusarum, Boeck, 1870,	{ 83, 140, 321, 337, 585 588, 1634, 1645
medusarum, Goës, 1865,	358
medusarum, ² Gosse, 1853,	275
minuta, Edward, 1868,	494, 588
oblivia, Krøyer, 1838,	{ 179, 189, 302, 306, 358 465, 494, 541, 588, 1419 1628, 1645
oblivia, Bate and Westwood,	{ 358, 382, 393, 541, 588 1419, 1628
pedestris, Guérin, 1836,	163
pelagica, M.-Edw., 1830,	102, 142, 175, 557
prehensilis, Bate and Westwood,	374, 494, 588
promontorii, Stebbing, 1888,	1385 , pl. clxvi, B
pupa, A. Costa, 1853,	274, 299 , 589, 1403
rubescens, Bovallius, 1887,	588
schizogeneios, Stebbing, 1888,	1391 , pl. clxviii
sibaginis, Stebbing, 1888,	1379 , pl. clxv
spinidorsalis, Sp. Bate, 1877,	473
spinipes, Boeck, 1860,	321, 358
{ Suerii, Latreille (in Desmarest),	122, 127, 137, 175, 189
1823-5,	
Sueurii, Gould, 1841,	194
tauriformis, Bate and Westwood,	374, 494, 558, 1628
1868,	
tricuspidata, Streets, 1877,	469, 588, 1671
trigona, Dana, 1852,	268, 589
Hyperiella—	
antarctica, Bovallius, 1887,	589, 592, 1407
dilatata, Stebbing, 1888,	1403 , pl. clxxi
fusca, Bovallius, 1887,	589
? pupa, Bovallius, 1887,	589
Hyperiopsis—	
Vøringii, G. O. Sars, 1885,	567, 572, 576

Hyperoche—

abyssorum, Bovallius, 1887,	588, 592
cryptodactylus, Stebbing, 1888,	1399 , pl. clxx
Krocyeri, Bovallius, 1887,	{ 588, 592, 594, 1399 1645
Luetkeni, Bovallius, 1887,	588, 592, 1399, 1645
Martinezi, Bovallius, 1887,	588, 1399
medusarum, Hansen, 1887,	592, 1399, 1628, 1645
prehensilis, Bovallius, 1887,	588
Iassa—	
pelagica, Leach, 1814,	86
pulchella, Leach, 1814,	86
Ichnopus—	
affinis, Heller, 1866,	366
calceolatus, Heller, 1866,	366, 545, 561
minutus, Boeck, 1870,	393
spinicornis, Boeck, 1860,	321, 361, 366, 591, 1650
taurus, Costa, 1853,	274, 296, 365, 545
umbonatus, G. O. Sars, 1882,	538
Ichthyomyzocus—	
Lophii, Hesse, 1873,	418, 464, 1630
Morrhue, Hesse, 1873,	418, 464, 1630
ornatus, Hesse, 1873,	418, 464, 1631
Squatinae, Hesse, 1873,	418, 464, 1631
Icilius—	
australis, Haswell, 1880,	511, 512, 533, 1208
dane, Stebbing, 1888,	1203 , pl. cxxxiii
ellipticus, Dana, 1852,	255, 265, 1208
ovalis, Dana, 1852,	255, 265
punctatus, Haswell, 1880,	512, 533
Icridium—	
fuscum, Grube, 1864,	{ 81, 341, 348, 354, 441 1627, 1638
Rissoanum, Catta, 1875,	81, 341, 441, 545
Iduna (see Liljeborgia)—	
brevicornis, Boeck, 1860,	322
fissicornis, Boeck, 1860,	322
Ione (also spelt Jone)—	
{ thoracica, Guérin-Ménéville, 1836,	163
thoracicus, Latreille, 1829,	137, 192
Iphigenia—	
abyssorum, Grimm, 1880,	509
Iphigenia—	
typica, G. M. Thomson, 1882,	586, 1637
Iphimedia—	
? ambigua, Haswell, 1880,	288, 512
capensis, Dana, 1852,	266, 928
carinata, Heller, 1866,	366
Eblanae, Sp. Bate, 1857,	295, 331, 348, 353, 434
fissicauda, Dana, 1852,	266, 447
minuta, G. O. Sars, 1882,	540
multispinis, Grube, 1864,	348, 353, 1627
nodosa, Dana, 1852,	255, 266, 353, 1627
Normani, Cunningham, 1870,	405
obesa, Rathke, 1843,	{ 205, 216, 219, 271, 295 305, 314, 331, 348, 366 395, 434, 442, 540, 545

¹ For the disentanglement, so far as practicable, of the synonymy of *Hyperia medusarum*, the papers by Bovallius and Hansen in 1887 should be consulted.

² Gosse's species, according to the Brit. Mus. Catal. Amph. Crust., pp. 292, 293, is *Hyperia galba*, Montagu.